

NOTISER

Errata & Addenda.

G. HENNINGSMOEN: "Remarks on the classification of trilobites" — Norsk geol. tidsskr., 29, pp. 174—217. Oslo 1951.

- p. 184, line 27: read *Xystridura* instead of *Xystridure*
p. 186, line 25, and p. 189, line 27: read *Bergeroniellus* instead of *Bergerionellus*
p. 193, line 13: read Nepeidae instead of Nereidae
p. 203, lines 28—29: read Conocoryphacea instead of Cococoryphacea
p. 204, line 16: read Swinnerton (1915) instead of Walcott (1891)
» line 19: read Olenellidea instead of Olenellidae
» line 20 and line 21: read (Swinnerton, 1915) instead of (Walcott, 1891)
p. 206, line 10: read Westergård, 1946 instead of Westergård, 1936
p. 208, line 13: read Nepeidae instead of Nereidae
p. 209, line 7: read Raphiophoracea instead of Raphiophoridaea
p. 205, line 28: The family name Cephalocoelidae Raymond, 1937 was changed to Catillicephalidae by Raymond, 1938 (Bull. Geol. Soc. Amer., suppl. to vol 48, p. XV), as *Cephalocoelia* was replaced by *Catillicephala*. It may be added here that the family Catillicephalidae, which was not assigned to any superfamily by Henningsmoen (1951, p. 205), has been shown by Shaw (1952, Jour. Paleont., 26, no. 3, p. 461) to belong to the Zacanthoidacea (= Corynexochida). Shaw (p. 466) has given good reasons for assuming that the cheirurid subfamily Sphaerexochinae has developed from the Catillicephalidae. This means that the superfamily Cheiruracea most probably developed from the Zacanthoidacea, not from the Conocoryphacea as suggested by Henningsmoen (1951, p. 203).
- p. 209: add *Alsataspidae* Turner, 1940 to the list of Raphiophoracean families. As *Alsataspis* is considered a synonym of *Seleneceme* by Whittington (1952, Jour. Paleont., 26, no. 1, p. 4) the name of the family was changed to Selenecemidae by him.

Paleontological Museum, Oslo.
August 1952.

G. Henningsmoen.